



Soils Map



State: **Minnesota**
 County: **Kandiyohi**
 Location: **6-119N-33W**
 Township: **Gennessee**
 Acres: **89.87**
 Date: **6/20/2017**



Maps Provided By:

 CUSTOMIZED ONLINE MAPPING
 © AgriData, Inc. 2017 www.AgriDataInc.com



Soils data provided by USDA and NRCS.

©2017 AgriData, Inc.

Area Symbol: MN067, Soil Area Version: 15

| Code | Soil Description | Acres | Percent of field | PI Legend | Non-Irr Class *c | Productivity Index | Bromegrass alfalfa | Bromegrass alfalfa hay | Corn | Oats | Reed canarygrass | Soybeans | Spring wheat |
|--------|--|-------|------------------|-----------|------------------|--------------------|--------------------|------------------------|------|------|------------------|----------|--------------|
| L191A | Blue Earth, Houghton, and Klossner soils, ponded, firm substratum, 0 to 1 percent slopes | 15.77 | 17.5% | | VIIIw | 5 | | | | | | | |
| L185B | Strout-Arkton complex, 2 to 6 percent slopes | 14.02 | 15.6% | | Ile | 83 | 5.1 | 4 | 152 | | | 46 | 49 |
| L184A | Corvuso silty clay loam, 0 to 2 percent slopes | 12.58 | 14.0% | | IIw | 78 | 4.6 | 3.6 | 143 | | | 43 | 48 |
| L186A | Danielson-Danielson, overwash complex, 1 to 4 percent slopes | 10.43 | 11.6% | | IIw | 93 | 5 | 3.8 | 170 | | | 51 | 49 |
| L166C2 | Newlondon-Strout complex, 6 to 12 percent slopes, moderately eroded | 9.67 | 10.8% | | IIIe | 77 | 4.5 | 3.4 | 141 | | | 42 | 44 |
| L164A | Lura silty clay, depressional, firm substratum, 0 to 1 percent slopes | 6.48 | 7.2% | | IIIw | 81 | 5 | 3.2 | 148 | | 5.5 | 45 | 47 |
| L166D2 | Newlondon-Strout complex, 12 to 18 percent slopes, moderately eroded | 5.74 | 6.4% | | IVe | 64 | 3.8 | 2.8 | 117 | | | 35 | 38 |
| 954D | Swanlake-Ves loams, 12 to 18 percent slopes | 5.62 | 6.3% | | IVe | 61 | 3.9 | 3 | 112 | 51 | | 34 | 30 |

| | | | | | | | | | | | | | |
|-------------------------|--|------|------|--|------|-------------|------------|------------|--------------|------------|------------|-------------|-------------|
| L33A | Kandiyohi clay, 0 to 2 percent slopes | 4.01 | 4.5% | | llw | 90 | 5 | 3.9 | 165 | | | 50 | 50 |
| 954C | Swanlake-Ves loams, 6 to 12 percent slopes | 3.22 | 3.6% | | llle | 78 | 5.1 | 3.9 | 142 | 75 | | 43 | 45 |
| L322A | Uniongrove loam, 0 to 2 percent slopes | 2.33 | 2.6% | | llw | 94 | 4.7 | 3.6 | 161 | | | 48 | 50 |
| Weighted Average | | | | | | 66.8 | 3.9 | 2.9 | 120.4 | 5.9 | 0.4 | 36.3 | 37.6 |

*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.